

β -Carbonic Anhydrase from *Pisum sativum*: Crystallization and Preliminary X-Ray Analysis

M. Kimber, J. Coleman, and E. Pai (U. of Toronto)

Abstract No. kimb9516

Beamline(s): X8C

Introduction: Carbonic anhydrases (CA) are enzymes that catalyze the reversible hydrolysis of CO₂ to bicarbonate. There are three different proteins with independent evolutionary origins, designated α -, β - and γ -CA. β -CA in plants plays a role as facilitator of high turnover in the photosynthetic dark reaction.

Methods and Materials: β -CA from garden peas was cloned and over expressed in *Escherichia coli* cells. Purified protein was crystallized by vapour phase equilibration against 16% PEG4000, 0.4 M ammonium acetate, 50 mM DTT and 100 mM sodium citrate pH 5.0. Macroseeding drastically improved the quality of the crystals. From a single flash-frozen crystal, a native data set complete to 2.8 Å resolution was collected at the NSLS, beamline X8C using radiation of 0.9500 Å wavelength and a MAR345 area detector. All data were reduced using the HKL package.

Results: Crystals of β -CA from garden peas adopted space group C222, unit cell parameters $a = 136.3$ Å, $b = 142.5$ Å, $c = 201.4$ Å. They grew to $0.2 \times 0.2 \times 0.08$ mm³ in size. Unit cell lengths, however, could vary up to 2% for different crystals. The crystals were also quite anisotropic in their scattering with diffraction limits varying between 2 Å and 4 Å. Self rotation analysis of the Patterson function revealed the presence of several weak two-fold rotation axes but no four-fold axis was found.

Acknowledgments: We would like to thank the staff at Brookhaven National Laboratory, beam line X8C (supported in part by a grant from MRC/NSERC), for their help and advice. We are also grateful to Patrick McGinn for purifying the initial batches of protein and Judith Jebanathirajah for help in maintaining cells. This research was supported by by NSERC and the MRC of Canada.